

SOLUTIONS

Math F251: Section 3.1 Worksheet

1. Differentiate the function.

$$F(r) = \frac{5}{r^3}$$

$$F(r) = 5 r^{-3}$$

$$F'(r) = -15 r^{-4}$$

$$y = 3e^x + \frac{4}{\sqrt[3]{x}}$$

$$y = 3e^x + 4x^{-\frac{1}{3}}$$

$$\frac{dy}{dx} = 3e^x - \frac{4}{3}x^{-\frac{4}{3}}$$

$$G(q) = (1 + q^{-1})^2$$

$$G(q) = 1^2 + 2q^{-1} + q^{-2} = 1 + 2q^{-1} + q^{-2}$$

$$G'(q) = -2q^{-2} - 2q^{-3}$$

2. Find equations of the tangent line and normal line to the curve at the given point: $y = x^2 + 2e^x$, $(0, 2)$

$$\frac{dy}{dx} = 2x + 2e^x$$

$$m_{\tan} = \left. \frac{dy}{dx} \right|_{x=0} = 0 + 2e^0 = 2, \quad m_{\text{normal}} = -\frac{1}{2}$$

Tangent line: $y - 2 = 2(x - 0)$

Normal line: $y - 2 = \left(-\frac{1}{2}\right)(x - 0)$

Math F251: Section 3.1 Worksheet

3. The equation of motion of a particle is $s = t^4 - 2t^3 + t^2 - t$, where s is in meters and t is in seconds.

(a) Find the velocity and acceleration as functions of t .

$$v = s' = \dot{s} = 4t^3 - 6t^2 + 2t - 1$$

$$a = v' = s'' = \ddot{s} = 12t^2 - 12t + 2$$

(b) Find the acceleration after 1 s.

$$a(1) = 12 - 12 + 2 = 2 \frac{m}{s^2}$$

4. Find an equation of a tangent line to the curve $y = x^4 + 1$ which is parallel to the line $32x - y = 15$.

line $y = 32x - 15$ has $m = 32$

$$\text{and } \frac{dy}{dx} = 4x^3 + 0 = 4x^3 \rightarrow \begin{array}{l} \text{solve} \\ 4x^3 = 32 \\ x = 2 \end{array}$$

$$y - 17 = 32(x - 2)$$

$$\Rightarrow \text{so } y = 2^4 + 1 = 17$$

5. Find the first and second derivatives of the function.

$$G(r) = \sqrt{r} - \sqrt[3]{r}$$

$$G(r) = r^{1/2} - r^{1/3}$$

$$G'(r) = \frac{1}{2}r^{-1/2} - \frac{1}{3}r^{-4/3}$$

$$G''(r) = -\frac{1}{4}r^{-3/2} + \frac{2}{9}r^{-5/3}$$