

Worksheet: Double and triple integrals!


1. Suppose $A = \{(x, y) \mid 1 \leq x^2 + y^2 \leq 4\}$. Write the double integral as an iterated integral, and evaluate it:

$$\iint_A \sqrt{x^2 + y^2} \, dA =$$

$x^2 + y^2 = r^2$
 $\sqrt{x^2 + y^2} = r$
 $dA = r \, dr \, d\theta$

(Hint. Sketch A. You can do the integral in polar coordinates!)

$$= \int_{\theta=0}^{2\pi} \int_{r=1}^2 r \cdot r \, dr \, d\theta$$



$$= 2\pi \int_1^2 r^2 \, dr = 2\pi \left[\frac{r^3}{3} \right]_1^2 = 2\pi \left(\frac{8}{3} - \frac{1}{3} \right)$$

$$= \left(\frac{14\pi}{3} \right)$$

2. The set $E = [0, 1] \times [1, 2] \times [2, 3]$ is a cube. Write the triple integral as an iterated integral, and evaluate it:

$$\iiint_E x + y \, dV = \int_{x=0}^1 \int_{y=1}^2 \int_{z=2}^3 x + y \, dz \, dy \, dx$$


$$= \int_0^1 \int_1^2 (x+y)(3-z) \, dy \, dx = \int_0^1 \left(\int_1^2 x + y \, dy \right) dx$$

$$= \int_0^1 \left[xy + \frac{y^2}{2} \right]_1^2 dx = \int_0^1 \left[2x + 2 - \left(x + \frac{1}{2} \right) \right] dx$$


$$= \int_0^1 x + \frac{3}{2} dx = \left[\frac{x^2}{2} + \frac{3}{2}x \right]_0^1 = \frac{1}{2} + \frac{3}{2} = \left(2 \right)$$

3. A right pyramid R has a base in the x, y plane which is the square $[-1, 1] \times [-1, 1]$, and its tip is at the point $(0, 0, 1)$. Its density increases as one approaches the tip, namely $\rho(x, y, z) = 1 + z$, in mass per volume units. Find the total mass.

$$M = \iiint_R (1+z) dV$$



$$= \int_{z=0}^1 \int_{x=z-1}^{x=1-z} \int_{y=z-1}^{y=1-z} (1+z) dy dx dz$$



CORRECTED

$$= \int_0^1 (1+z) (\text{area of square}) dz$$

$$= \int_0^1 (1+z) ((z-1, 1-z) \times (z-1, 1-z)) dz$$

$$= \int_0^1 (1+z) ((1-z - z+1)^2) dz = 2 \int_0^1 (1+z)(1-z)^2 dz = 4 \int_0^1 (1-z^2)(1-z) dz$$

4. Find the volume of the sphere of radius one by setting-up either a double or a triple integral, and evaluating it. Of course, the answer you should get is $V = 4\pi/3$.

$$V = 2 \int_{\theta=0}^{2\pi} \int_{r=0}^1 \sqrt{1-r^2} r dr d\theta = 4\pi \int_0^1 (1-r^2)^{1/2} r dr$$

$$= 4 \int_0^1 (1-z-z^2) dz$$

$$= 4 \left(1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} \right)$$

$$= \frac{5}{3}$$

$$= 4\pi \int_1^0 u^{1/2} \frac{du}{-2} = 2\pi \int_0^1 u^{1/2} du = 2\pi \left[\frac{2}{3} u^{3/2} \right]_0^1 = \frac{4\pi}{3}$$

$$V = \int_{x=-1}^1 \int_{y=-\sqrt{1-x^2}}^{y=\sqrt{1-x^2}} \int_{z=-\sqrt{1-x^2-y^2}}^{z=\sqrt{1-x^2-y^2}} 1 dz dy dx$$

wise to switch to polar here

$$= \int_{-1}^1 \int_{y=-\sqrt{1-x^2}}^{y=\sqrt{1-x^2}} 2\sqrt{1-x^2-y^2} dy dx = \int_{\theta=0}^{2\pi} \int_{r=0}^1 2\sqrt{1-r^2} r dr d\theta$$

$$= \dots = \frac{4\pi}{3}$$