2.2 Separable Equations a lecture for MATH F302 Differential Equations

Ed Bueler, Dept. of Mathematics and Statistics, UAF

Fall 2023

for textbook: D. Zill, A First Course in Differential Equations with Modeling Applications, 11th ed.

regarding chapters 1 and 2

- the purpose of textbook chapter 1 is to sketch the language and meaning of differential equations
- the purpose of chapter 2 is to actually solve some differential equations by hand
 - section 2.2 is the first example: separable equations
 - warning: by-hand methods can be difficult or impossible on a given differential equation
 - however, the examples where we know how to solve are often important in practice

separable differential equations

Definition

a separable differential equation can be put in the form

$$\frac{dy}{dx} = g(x)h(y)$$

• Example.

$$\frac{dy}{dx} = \frac{y\cos(x)}{1+y^2}$$

• Not an example.

$$\frac{dy}{dx} = \cos(x) + y$$

• Also **not** an example.

$$\frac{dy}{dx} = \sin(x + y^2)$$

emphasis on can

Definition

a separable differential equation can be put in the form

$$\frac{dy}{dx} = g(x)h(y)$$

• Example.

$$p(y)\frac{dy}{dx} = g(x)$$

emphasis on can

Definition

a separable differential equation can be put in the form

$$\frac{dy}{dx} = g(x)h(y)$$

• Example.

$$p(y)\frac{dy}{dx} = g(x)$$

define h(y) = 1/p(y) to make into standard separable form

• move y stuff to left and x to right:

$$\frac{dy}{dx} = g(x)h(y)$$
$$\frac{1}{h(y)}\frac{dy}{dx} = g(x)$$
$$\frac{1}{h(y)}dy = g(x) dx$$

• move y stuff to left and x to right:

$$\frac{dy}{dx} = g(x)h(y)$$
$$\frac{1}{h(y)}\frac{dy}{dx} = g(x)$$
$$\frac{1}{h(y)}dy = g(x) dx$$

• integrate both sides:

$$\int \frac{1}{h(y)} \, dy = \int g(x) \, dx$$

• alternative appearance with p(y) = 1/h(y):

$$p(y)\frac{dy}{dx} = g(x)$$

• alternative appearance with p(y) = 1/h(y):

$$p(y)\frac{dy}{dx} = g(x)$$

• move y stuff to left and x to right:

$$p(y)\,dy=g(x)\,dx$$

• integrate both sides:

$$\int p(y)\,dy = \int g(x)\,dx$$

why does it work?

- the method works because of the chain rule
- the integrals you are really doing are both with respect to x:

$$\int p(y(x)) \frac{dy}{dx} \, dx = \int g(x) \, dx$$

how do you finish up?

once you do the integrals

$$\int p(y)\,dy = \int g(x)\,dx$$

then solve for y, if possible, to get an explicit solutionif you cannot solve for y then the solution remains implicit

• example: find y(x) if

$$\frac{dy}{dx} = xy^2$$

- some familiar equations are also separable
- example: find y(x) if

$$\frac{dy}{dx} = -5y$$

- what if there are initial conditions?
- example: find z(t) if z(4) = 1 and $z' = \frac{e^{-z}}{t}$

• you may end up only knowing the solution implicitly

• example: find
$$y(x)$$
 if $\frac{dy}{dx} = \frac{x(1-x)}{y(2+y)}$

- integration techniques are required, and
- division is dangerous!
- example: find all solutions y(x) if

$$\frac{dy}{dx} = y^2 - 9$$

standard expectations

to learn this material, just listening to a lecture is not enough

- please read section 2.2 in the textbook
- please *do* the Homework for section 2.2
- search "separable ODEs" at YouTube to see more examples