

# SOLUTIONS

## Von Neumann analysis: plug and chug

Corrected

Section 9.6 of the textbook (R. J. LeVeque, 2007) presents von Neumann analysis without emphasizing how people actually do the analysis. This worksheet exercises the standard style.<sup>1</sup>

**Example. FTCS on heat equation.** It is easiest to explain the idea relative to an example. Suppose we apply the FTCS scheme to the heat equation  $u_t = Du_{xx}$  with constant diffusivity  $D > 0$ :

$$(1) \quad \frac{U_j^{n+1} - U_j^n}{k} = D \frac{U_{j-1}^n - 2U_j^n + U_{j+1}^n}{h^2}$$

To find what time steps  $k > 0$  would be stable for a given spacing  $h > 0$ , von Neumann substituted

$$(2) \quad U_j^n = g(\xi)^n e^{ijh\xi}$$

into scheme (1), where  $\xi \in \mathbb{R}$  is the wave number for the spatial wave  $e^{ijh\xi}$ , in which  $i = \sqrt{-1} \in \mathbb{C}$  (as usual). The scalar function  $g(\xi)$  is called the amplification factor of the scheme.

The spatial wave is complex, but it really is a wave. In fact, for the interval  $0 \leq x \leq 1$  the grid is  $x_j = jh$  and thus

$$e^{ijh\xi} = e^{i\xi x_j} = \cos(\xi x_j) + i \sin(\xi x_j).$$

Now we want to find  $g(\xi)$ . To compute it, substitute form (2) into scheme (1). Indices “ $n+1$ ,” “ $j-1$ ,” and “ $j+1$ ” will turn into powers. Then use the properties of the exponential. After simplification and trigonometric identities—do the details in Exercise 1—we get


$$g(\xi) = 1 - \frac{4Dk}{h^2} \sin^2\left(\frac{\xi h}{2}\right).$$

Absolute stability  $|U_j^{n+1}| \leq |U_j^n|$  corresponds to  $|g(\xi)| \leq 1$  for all  $\xi \in \mathbb{R}$ . For this scheme<sup>2</sup> we get the condition  $k \leq \frac{h^2}{2D}$ , which tells us that the time step  $k$  must be very small if the spacing  $h$  is small.

**Exercise 1. FTCS on heat equation.** Label the stencil. Then fill in the above details.

$[u_t = Du_{xx} \Rightarrow (1)]$

put  $U_j^n = g(\xi)^n e^{ijh\xi}$  into (1):



$$\frac{g(\xi)^{n+1} e^{ijh\xi} - g(\xi)^n e^{ijh\xi}}{k} = D \frac{g(\xi)^n e^{i(j-1)h\xi} - 2g(\xi)^n e^{ijh\xi} + g(\xi)^n e^{i(j+1)h\xi}}{h^2}$$

$$D \frac{g(\xi)^n e^{i(j-1)h\xi} - 2g(\xi)^n e^{ijh\xi} + g(\xi)^n e^{i(j+1)h\xi}}{h^2}$$

now cancel  $g(\xi)$  &  $e^{ijh\xi}$ :

$$\frac{g(\xi) - 1}{k} = D \frac{e^{-ih\xi} - 2 + e^{ih\xi}}{h^2} \Rightarrow g(\xi) = 1 + R(e^{-ih\xi} + e^{ih\xi} - 2)$$

$$\begin{aligned} R &= \frac{Dk}{h^2} \rightarrow R(e^{-ih\xi} + e^{ih\xi} - 2) \\ &= 1 + R(2\cos(h\xi) - 2) \\ &= 1 - 2R(1 - \cos(h\xi)) \end{aligned}$$

<sup>1</sup>The textbook eventually says it clearly in section 10.5.

<sup>2</sup>The same condition is derived via MOL in section 9.3, and then by von Neumann analysis in section 9.6.

for all  $\xi$


$$\begin{aligned} |g(\xi)| \leq 1 &\Leftrightarrow 1 - 4R \sin^2(h\xi/2) \geq -1 \\ &\Leftrightarrow 4R \sin^2(h\xi/2) \leq 2 \Rightarrow k \leq \frac{h^2}{2D} \end{aligned}$$

$$\begin{aligned} &= 1 - 4R \cdot \frac{1}{2} (1 - \cos(h\xi)) \\ &= 1 - 4R \sin^2(h\xi/2) \end{aligned}$$

I did not give you enough space! Only highlights are shown.

Exercise 2. Crank-Nicolson on heat equation. Label the stencil. State the scheme. Do the analysis.

put  $U_j^n = g(\xi)^n e^{ijh\xi}$  into CN scheme and cancel:



$$\frac{g(\xi) - 1}{k} = \frac{D}{2h^2} (g(\xi) e^{-i h \xi} - 2g(\xi) + g(\xi) e^{i h \xi} + e^{-i h \xi} - 2 + e^{i h \xi})$$

rearrange, use  $\cos \theta = \frac{1}{2}(e^{-i\theta} + e^{i\theta})$ , use  $R = Dk/h^2$ :

$$(R - R \cos(h\xi) + 1) g(\xi) = 1 + R \cos(h\xi) - R$$

rearrange use  $\sin^2 \theta = \frac{1}{2}(1 - \cos(2\theta))$ :


$$g(\xi) = \frac{1 - 2R \sin^2(h\xi/2)}{1 + 2R \sin^2(h\xi/2)} \Rightarrow |g(\xi)| \leq 1 \Rightarrow \text{for all } \xi$$

stable for any  $k > 0$

CORRECTED

Exercise 3. FTCS on the advection equation  $u_t + au_x = 0$  with  $a$  constant. Write down the scheme, draw and label a stencil, and do the analysis.

scheme: 
$$\frac{U_j^{n+1} - U_j^n}{k} + a \frac{U_{j+1}^n - U_{j-1}^n}{2h} = 0$$



put  $U_j^n = g(\xi)^n e^{ijh\xi}$  into scheme, cancel, use  $r = ak/h$ :

$$g(\xi) - 1 + \frac{1}{2} r (e^{i h \xi} - e^{-i h \xi}) = 0$$

use  $\sin \theta = (e^{i\theta} - e^{-i\theta})/2i$ , rearrange:

$$g(\xi) = 1 - i r \sin(h\xi)$$

magnitude:  $|g(\xi)| = \sqrt{1 + r^2 \sin^2(h\xi)}$

$|g(\xi)| > 1$  for  $k > 0, \xi \neq 0$

not stable for any  $k$

Exercise 4. CTCS (leapfrog) on advection equation. Again.

scheme: 
$$\frac{U_j^{n+1} - U_j^{n-1}}{2k} + a \frac{U_{j+1}^n - U_{j-1}^n}{2h} = 0$$

put  $U_j^n = \dots$  into scheme, cancel, use  $r = ak/h$ :

$$g(\xi) - g(\xi)^{-1} + r (e^{i h \xi} - e^{-i h \xi}) = 0$$

use  $\sin \theta = (e^{i\theta} - e^{-i\theta})/2i$ , rearrange:

$$g(\xi)^2 + 2i r \sin(h\xi) g(\xi) - 1 = 0$$

if  $|r| \leq 1$  then  $|g(\xi)| = 1$

so stable if  $|ak/h| \leq 1$

quad. formula: 
$$g(\xi) = -i r \sin(h\xi) \pm \sqrt{1 - r^2 \sin^2(h\xi)}$$