

**Problem 14.** Suppose  $A, B$  are disjoint sets with  $A \cup B = \mathbb{R}$ , and suppose that  $a < b$  for all  $a \in A$  and  $b \in B$ . Then there exists  $c \in \mathbb{R}$  such that  $x \leq c$  for  $x \in A$  and  $x \geq c$  for  $x \in B$ .

*Proof.* □

**Problem 15.** Here is an example which shows that the claim in Problem 14 is false if  $\mathbb{R}$  is replaced, in both instances, by the set of rationals  $\mathbb{Q}$ :

**Problem 16.** Let  $a < b$  be real numbers. Define the set  $T = \mathbb{Q} \cap [a, b]$ . Then  $\sup T = b$ .

*Proof.* □

**Problem 17.** By definition, a set  $C \subseteq \mathbb{R}$  is dense if for any real numbers  $a < b$  there is  $c \in C$  so that  $a < c < b$ . Let  $T$  be the set of all rational numbers  $p/q$ , with  $p \in \mathbb{Z}$ , for which  $q = 2^k$  for some  $k \in \mathbb{N}$ . Then  $T$  is dense.

*Proof.* □

**Problem 18.**

- (a) An example of two real sets  $A, B$  with  $A \cap B = \emptyset$ ,  $\sup A = \sup B$ ,  $\sup A \notin A$ , and  $\sup B \notin B$  is
- (b) An example of a sequence of nested open intervals  $J_1 \supseteq J_2 \supseteq J_3 \supseteq \dots$ , with  $S = \bigcap_{n=1}^{\infty} J_n$  nonempty and of finite cardinality, is
- (c) By definition, an unbounded closed interval is of the form  $[a, \infty) = \{x \in \mathbb{R} : x \geq a\}$ . An example of a sequence of nested unbounded closed intervals  $L_1 \supseteq L_2 \supseteq L_3 \supseteq \dots$ , with  $\bigcap_{n=1}^{\infty} L_n = \emptyset$ , is

**Problem 19.** If  $A \subseteq B$  and  $B$  is countable then  $A$  is either countable or finite.

*Proof.* Assume  $B$  is countable. If  $|A| < \infty$  then  $A$  is finite and we are done. So we will consider an infinite subset  $A \subseteq B$  and show it is countable. □

**Problem 20.**

- (a) For any  $a < b$  it follows that  $(a, b) \sim \mathbb{R}$ .

*Proof.* □

- (b)  $[0, 1) \sim (0, 1)$

*Proof.* □

**Problem 21.** If  $A \sim B$  and  $B \sim C$  then  $A \sim C$ .

*Proof.* □